

CHEMISTRY 101
Hour Exam II
March 14, 2019
McCarren

Name _____

Signature _____

Section _____

This exam contains 17 questions on 10 numbered pages. Check now to make sure you have a complete exam. You have one hour and thirty minutes to complete the exam. Determine the best answer to the first 15 questions and enter these on the special answer sheet. Also, circle your responses in this exam booklet. Show all of your work and provide complete answers to questions 16 and 17. A periodic table and one sheet of scratch paper are provided after the exam. Anything written on the periodic table and scratch paper will not be graded.



1-15	(30 pts.)	_____
16	(12 pts.)	_____
17	(18 pts.)	_____
Total	(60 pts.)	_____

Useful Information:

1 L = 1000 mL (exactly)

Always assume ideal behavior for gases (unless explicitly told otherwise).

$PV = nRT$

$R = 0.08206 \text{ L} \cdot \text{atm/mol} \cdot \text{K} \approx 0.0821 \text{ L} \cdot \text{atm/mol} \cdot \text{K}$

$K = ^\circ\text{C} + 273$


$N_A = 6.022 \times 10^{23} = 1 \text{ mole}$

Solubility Rules:

1. Most nitrate salts are soluble.
2. Most salts of sodium, potassium, and ammonium cations are soluble.
3. Most chloride salts are soluble. Exceptions: silver(I), lead(II), and mercury(I) chloride.
4. Most sulfate salts are soluble. Exceptions: calcium, barium, and lead(II) sulfate.
5. Most hydroxide salts can be considered insoluble. Soluble ones: sodium, potassium, ammonium, and calcium hydroxide.
6. Consider sulfide, carbonate, and phosphate salts to be insoluble. Soluble ones: sodium, potassium, and ammonium.






Part 1: Multiple Choice

1. Balance the equation shown below. Which image shows the correct particle representation of the equation after it has been balanced? Assume that A and B each represent atoms of different elements.




Key:



a.	
b.	
c.	
d.	
e.	

2. The balanced equation for the reaction between hydrogen and phosphorus to make phosphine gas is shown below. Which statement is **always true** about this reaction?




- Hydrogen gas is limiting because it is used up twice as fast as phosphorus.
- Every time three molecules of hydrogen gas are used, two molecules of phosphine form.
- The reaction container holds four moles of phosphine gas after the reaction.
- Before a reaction starts, there must be at least 3 moles of phosphorus and 6 moles of hydrogen present.
- At least two of these statements are always true about this reaction.

Consider the reaction between solid diphosphorus pentoxide and liquid water to form aqueous phosphoric acid (H_3PO_4). Write out and balance the equation. Then, use this equation to answer the next three questions.

3. Balance the equation and determine the sum of the coefficients when the equation is balanced in standard form (lowest whole number ratios).
 - a. 3
 - b. 5
 - c. 6
 - d. 9
 - e. 12
4. If 6.0 moles of phosphoric acid were formed as a result of this reaction, how many moles of water were needed to react?
 - a. 1.0 mole
 - b. 3.0 moles
 - c. 6.0 moles
 - d. 9.0 moles
 - e. 12 moles
5. If 6.0 moles of phosphoric acid were formed in this reaction, what mass of diphosphorus pentoxide was needed to react?
 - a. 70.95 g
 - b. 141.9 g
 - c. 283.8 g
 - d. 425.7 g
 - e. 851.4 g

Please go on to the next page.

Consider the four beakers below, each containing an aqueous salt dissolved in water as shown. Aqueous iron(III) nitrate has been added to beakers 1 and 2. Use this information to answer the next three questions.



6. Addition iron(III) nitrate to beaker 1 containing aqueous potassium chromate resulted in the formation of a precipitate. What is the formula of the precipitate that has formed in beaker 1?
- $\text{Fe}_2(\text{CrO}_4)_3$
 - FeCrO_4
 - $\text{Fe}_2(\text{CO}_3)_3$
 - FeCO_3
 - KNO_3
7. When iron(III) nitrate was added to the aqueous salt solution in beaker 2, no precipitate formed. Which of the following salts may have been present in beaker 2?
- K_2SO_4
 - K_2CO_3
 - KOH
 - K_3PO_4
 - At least two of these salts may have been the salt present in beaker 2.
8. When the iron(III) nitrate solution is poured into beakers 3 and 4, will any precipitates form?
- A precipitate will form in beaker 3 only.
 - A precipitate will form in beaker 4 only.
 - Precipitates will form in both beakers 3 and 4.
 - Precipitates will form in neither beakers 3 nor 4.
 - More information is needed to determine whether or not a precipitate will form in either of these beakers.

An aqueous solution of calcium chloride reacts with an aqueous solution of sodium phosphate. Use this reaction to answer the next several questions.

9. What is the **complete** ionic equation for this reaction?

- a. $3\text{Ca}^{+2}(\text{aq}) + 6\text{Cl}^{-}(\text{aq}) + 6\text{Na}^{+}(\text{aq}) + 2\text{PO}_4^{3-}(\text{aq}) \rightarrow \text{Ca}_3(\text{PO}_4)_2(\text{s}) + 6\text{Na}^{+}(\text{aq}) + 6\text{Cl}^{-}(\text{aq})$
- b. $3\text{Ca}^{+2}(\text{aq}) + 6\text{Cl}_2^{-}(\text{aq}) + 6\text{Na}_3^{+}(\text{aq}) + 2\text{PO}_4^{3-}(\text{aq}) \rightarrow 3\text{Ca}^{+2}(\text{aq}) + 2\text{PO}_4^{3-}(\text{aq}) + 6\text{Na}_3\text{Cl}_2(\text{s})$
- c. $3\text{Ca}^{+2}(\text{aq}) + 6\text{Cl}^{-}(\text{aq}) + 6\text{Na}^{+}(\text{aq}) + 2\text{PO}_4^{3-}(\text{aq}) \rightarrow 3\text{Ca}^{+2}(\text{aq}) + 2\text{PO}_4^{3-}(\text{aq}) + 6\text{NaCl}(\text{s})$
- d. $\text{Ca}^{+2}(\text{aq}) + \text{Cl}^{-}(\text{aq}) + \text{Na}^{+}(\text{aq}) + \text{PO}_4^{3-}(\text{aq}) \rightarrow \text{CaPO}_4(\text{s}) + \text{Na}^{+}(\text{aq}) + \text{Cl}^{-}(\text{aq})$
- e. $2\text{Ca}^{+2}(\text{aq}) + 6\text{Cl}^{-}(\text{aq}) + 6\text{Na}^{+}(\text{aq}) + 3\text{PO}_4^{3-}(\text{aq}) \rightarrow \text{Ca}_2(\text{PO}_4)_3(\text{s}) + 6\text{Na}^{+}(\text{aq}) + 6\text{Cl}^{-}(\text{aq})$

10. What is the **net ionic** equation for this reaction?

- a. $2\text{Cl}^{-}(\text{aq}) + 3\text{Na}^{+}(\text{aq}) \rightarrow \text{Na}_3\text{Cl}_2(\text{s})$
- b. $\text{Cl}^{-}(\text{aq}) + \text{Na}^{+}(\text{aq}) \rightarrow \text{NaCl}(\text{s})$
- c. $\text{Ca}^{+2}(\text{aq}) + \text{PO}_4^{3-}(\text{aq}) \rightarrow \text{CaPO}_4(\text{s})$
- d. $2\text{Ca}^{+2}(\text{aq}) + 3\text{PO}_4^{3-}(\text{aq}) \rightarrow \text{Ca}_2(\text{PO}_4)_3(\text{s})$
- e. $3\text{Ca}^{+2}(\text{aq}) + 2\text{PO}_4^{3-}(\text{aq}) \rightarrow \text{Ca}_3(\text{PO}_4)_2(\text{s})$

The solution of calcium chloride had a volume of 2.00 L and a concentration of 3.5 M. The sodium phosphate solution that it was mixed with had a volume of 2.00 L and a concentration of 2.0 M. Use this information as well as your equations above to answer the next two questions.


11. Which ions were present in the container after the reaction?

- a. Na^{+} and Cl^{-} only
- b. PO_4^{3-} and Cl^{-} only
- c. Na^{+} , PO_4^{3-} , and Cl^{-}
- d. Na^{+} , Ca^{+2} , and Cl^{-}
- e. Na^{+} , PO_4^{3-} , Ca^{+2} , and Cl^{-}

12. What is the concentration of **sodium** ions present in the container **after** the reaction? If these were not present in the container after the reaction, select 0 M.

- a. 0 M
- b. 1.0 M
- c. 2.0 M
- d. 3.0 M
- e. 6.0 M

Recall the lab experiment in which you observed several balloons inflating after reacting two different acids with sodium bicarbonate (baking soda). One of the reactions you saw took place below between the baking soda and sulfuric acid (H_2SO_4).



You have a flask containing sulfuric acid with a balloon holding baking soda on top of it. The baking soda is poured into the flask, the reaction occurs, and the balloon inflates. In the container, 42.0 grams of baking soda (molar mass 84.0 g/mol) combined with 500.0 mL of 1.00 M sulfuric acid, and the balloon filled with carbon dioxide at 1.10 atm and 25.0°C. Use this information to answer the next two questions.

13. What volume of carbon dioxide gas was produced in this reaction?
 - a. 2.78 L
 - b. 5.56 L
 - c. 11.1 L
 - d. 16.7 L
 - e. 22.2 L

14. How many moles of excess reactant were leftover after this reaction?
 - a. 0.125 moles
 - b. 0.250 moles
 - c. 0.500 moles
 - d. 0.750 moles
 - e. 1.00 moles

15. A second balloon contains 15.0 grams of baking soda reacting with 500.0 mL of 1.00 M sulfuric acid. These also react and fill the balloon with carbon dioxide. Will this balloon be larger, smaller, or equal to the size of the balloon in questions 12 and 13?
 - a. Larger: More carbon dioxide is produced, so the balloon is larger.
 - b. Larger: Due to less baking soda in the container, there is more room for the gas to expand, so the balloon is larger.
 - c. The same size: The same amount of sulfuric acid is used up, so the same amount of carbon dioxide is produced and the balloons are the same size.
 - d. Smaller: Less carbon dioxide is produced because the sulfuric acid is limiting, so the balloon is smaller.
 - e. Smaller: Less carbon dioxide is produced because the baking soda is limiting, so the balloon is smaller.

Part 2: Free Response

Please write in your answers completely, showing all work and explaining as necessary.

16. Consider each of the pairs of solutions below and use them to answer question 16.

Set 1

- **Solution A** consists of 60.0 grams sodium hydroxide dissolved to make 500. mL of solution.
- **Solution B** consists of 0.25 moles sodium hydroxide dissolved to make 250. mL of solution.

a. Which solution has a greater overall molarity? Show work and/or explain your answer.

b. For the solution in part a. that you said has the greater concentration, what volume of water (in mL) would you need to add to that solution to result in both solutions A and B having the same concentration? Show work.

Please go on to the next page.

Set 2


- **Solution C** consists of a 1.0 L aqueous solution of calcium hydroxide.
- **Solution D** consists of a 1.0 L aqueous solution of hydrochloric acid (HCl)

Solutions C and D are combined together.

- c. Give the balanced molecular equation for this reaction, showing all phases.
- d. When these solutions reacted, they reacted in a perfectly stoichiometric ratio; neither reactant was in excess and both reactants ran out at the same time. Which solution had a higher concentration, solution C or D? Explain your answer.

Please go on to the next page.

17. Consider the unbalanced equation between some unknown element X and fluorine gas (F_2) to form some compound XF_n containing both X and fluorine, where n represents some unknown whole number.



Before the reaction, six moles of X and six moles of fluorine gas are present in a sealed container. These later react to form the XF_n compound.

After the reaction:

- All six moles of fluorine gas have been consumed.
 - Four moles of element X remain unreacted.
 - Two moles of XF_n product have been formed.
- a. Even without having a balanced equation, this is enough information to complete a BCA table for the reaction. Use the information above to completely fill out each blank in the BCA table below:

	X	+	F_2	\rightarrow	XF_n
B	_____		_____		_____
C	_____		_____		_____
A	_____		_____		_____

- b. Using information from your BCA table, give the lowest whole number coefficients for each of the substances in this reaction. Explain how your BCA table displays this information.

Coefficient of X: _____ Coefficient of F_2 : _____ Coefficient of XF_n : _____

- c. What is the value of “n” in XF_n ? Show work and/or explain.

- d. What was the mass of the six moles of F_2 present before the reaction?
- e. If the total mass in the container before the reaction was 1,016 total grams of X and F_2 combined, what was the total mass of the six moles of X present?
- f. Use your answer to part e. to determine the identity of X.
- g. We have seen in chemical reactions that mass is conserved and therefore is not created or destroyed as a result of a chemical reaction. You now have all the information you need to demonstrate that mass was conserved in this reaction. To do this, fill in the table below by giving the masses of X, F_2 , and XF_n present before and after the reaction as well as the total mass after the reaction. Show all work in the space below the table.

	X	F_2	XF_n product	Total
Before reaction			0 grams	1,016 grams
After reaction				



This is the end of the exam. Nothing written after this page will be graded.

Chem 101 Scratch Paper

NOTHING WRITTEN ON THIS PAGE WILL BE GRADED

1A

1
H
Hydrogen
1.008

2A

3
Li
Lithium
6.941

4
Be
Beryllium
9.012

5
B
Boron
10.81

6
C
Carbon
12.01

7
N
Nitrogen
14.01

8
O
Oxygen
16.00

9
F
Fluorine
19.00

10
Ne
Neon
20.18

3A

11
Na
Sodium
22.99

12
Mg
Magnesium
24.31

13
Al
Aluminum
26.98

14
Si
Silicon
28.09

15
P
Phosphorus
30.97

16
S
Sulfur
32.07

17
Cl
Chlorine
35.45

18
Ar
Argon
39.95

19
K
Potassium
39.10

20
Ca
Calcium
40.08

21
Sc
Scandium
44.96

22
Ti
Titanium
47.88

23
V
Vanadium
50.94

24
Cr
Chromium
52.00

25
Mn
Manganese
54.94

26
Fe
Iron
55.85

27
Co
Cobalt
58.93

28
Ni
Nickel
58.69

29
Cu
Copper
63.55

30
Zn
Zinc
65.38

31
Ga
Gallium
69.72

32
Ge
Germanium
72.59

33
As
Arsenic
74.92

34
Se
Selenium
78.96

35
Br
Bromine
79.90

36
Kr
Krypton
83.80

37
Rb
Rubidium
85.47

38
Sr
Strontium
87.62

39
Y
Yttrium
88.91

40
Zr
Zirconium
91.22

41
Nb
Niobium
92.91

42
Mo
Molybdenum
95.94

43
Tc
Technetium
(98)

44
Ru
Ruthenium
101.1

45
Rh
Rhodium
102.9

46
Pd
Palladium
106.4

47
Ag
Silver
107.9

48
Cd
Cadmium
112.4

49
In
Indium
114.8

50
Sn
Tin
118.7

51
Sb
Antimony
121.8

52
Te
Tellurium
127.6

53
I
Iodine
126.9

54
Xe
Xenon
131.3

55
Cs
Cesium
132.90

56
Ba
Barium
137.3

57
La
Lanthanum
138.9

58
Ce
Cerium
140.12

59
Pr
Praseodymium
140.91

60
Nd
Neodymium
144.24

61
Pm
Promethium
(144.91)

62
Sm
Samarium
150.36

63
Eu
Europium
151.96

64
Gd
Gadolinium
157.25

65
Tb
Terbium
158.93

66
Dy
Dysprosium
162.50

67
Ho
Holmium
164.93

68
Er
Erbium
167.26

69
Tm
Thulium
168.93

70
Yb
Ytterbium
173.05

71
Lu
Lutetium
174.97

72
Hf
Hafnium
178.5

73
Ta
Tantalum
180.9

74
W
Tungsten
183.9

75
Re
Rhenium
186.2

76
Os
Osmium
190.2

77
Ir
Iridium
192.2

78
Pt
Platinum
195.1

79
Au
Gold
197.0

80
Hg
Mercury
200.6

81
Tl
Thallium
204.4

82
Pb
Lead
207.2

83
Bi
Bismuth
209.0

84
Po
Polonium
(209)

85
At
Astatine
(210)

86
Rn
Radon
(222)

87
Fr
Francium
(223)

88
Ra
Radium
226

89
Ac
Actinium
(227)

90
Th
Thorium
232.04

91
Pa
Protactinium
231.04

92
U
Uranium
238.03

93
Np
Neptunium
237.05

94
Pu
Plutonium
244.06

95
Am
Americium
243.06

96
Cm
Curium
247.07

97
Bk
Berkelium
247.07

98
Cf
Californium
251.08

99
Es
Einsteinium
252.08

100
Fm
Fermium
257.10

101
Md
Mendelevium
258.10

102
No
Nobelium
259.10

103
Lr
Lawrencium
262.10

104
Rf
Rutherfordium
(261)

105
Db
Dubnium
(262)

106
Sg
Seaborgium
(263)

107
Bh
Bohrium
(262)

108
Hs
Hassium
(265)

109
Mt
Meitnerium
(266)

110
Ds
Darmstadtium
(269)

111
—
(272)

112
—
(277)

113
Nh
Nihonium
(284)

114
—
(285)

115
Mc
Moscovium
(288)

116
—
(289)

117
Ts
Tennessine
(294)

118
Og
Oganesson
(294)

Key

Atomic number → Symbol

Name → Atomic mass

→ 67 Ho →

8A

1
He
Helium
4.003

2
Ne
Neon
20.18

3
Ar
Argon
39.95

4
Kr
Krypton
83.80

5
Xe
Xenon
131.3

6
Rn
Radon
(222)

7

Lanthanides

Actinides